A Pre-PhD Synopsis

"Studies on Bacterial Pathogens and their effects on Physiological and Nutritional Status of *Hypophthalmichthys molitrix* from Limboti Dam, District Nanded, M.S."

Submitted to

Swami Ramanand Teerth Marathwada University, Nanded

For

Research Registration Committee

Under

Faculty of Science and Technology

By

Baldawa Mohini Rajkumar

Under Guidance of

Dr. R. D. Barde

Professor & Head,

Department of Zoology,
Shri Guru Buddhiswami Mahavidhyalaya, Purna (Jn.)
Tq. Purna, Dist. Parbhani-431511

May 2025

"Studies n Bacterial Pathogens and their effects on Physiological and Nutritional status of *Hypophthalmichthys molitrix* from Limboti Dam, District Nanded, M.S."

Research Scholar: Baldawa Mohini Rajkumar

Research Supervisor: Dr. R. D. Barde

Abstract:

Hypophthalmichthys molitrix, an ecologically significant and nutritionally valuable filter-feeding species, plays a critical role in India's inland aquaculture systems, particularly in semi-natural reservoirs such as Limboti Dam, Ahmedpur. However, the increasing susceptibility of *Hypophthalmichthys molitrix* to bacterial infections is linked to the intensification of aquaculture practices and deteriorating water quality due to anthropogenic influences. Pathogens like *Aeromonas hydrophila*, *Pseudomonas fluorescens*, and *Edwardsiella tarda* release virulence enzymes—proteases, lipases, and lecithinases—which impair host immunity, resulting in tissue damage, physiological dysfunction, and nutritional imbalance.

This study aims to isolate and characterize the bacterial pathogens affecting silver carp in Limboti Dam and to assess their virulence potential via enzymatic profiling. A multidisciplinary approach will be employed, including haematological evaluations, histopathological examination of vital organs (gills, liver, kidney), and nutritional analysis of muscle tissues (protein, lipid, carbohydrate, and vitamin content). Additionally, environmental parameters such as water temperature, pH, salinity, dissolved oxygen, and ammonia will be monitored to understand their role in modulating disease severity and pathogen dynamics.

The findings are expected to bridge a critical gap in reservoir-based fish health management by establishing statistically significant relationships between bacterial burden, virulence expression, and the physiological and nutritional degradation of *Hypophthalmichthys molitrix*. The outcomes will aid in the development of early disease detection systems, contribute baseline epidemiological data, and inform sustainable disease mitigation strategies tailored to inland aquaculture in Maharashtra.

Keywords:*Hypophthalmichthys molitrix*, bacterial pathogens, physiological stress, nutritional depletion, inland aquaculture management.

1. Introduction

Emerging as a major part of world food systems, freshwater aquaculture offers a sustainable source of protein to satisfy the nutritional needs of rising population (Little et al., 2016). Due in

great part to their high growth rate, filter-feeding behavior, and adaptation to pond and reservoir conditions, freshwater fish—especially species of the Indian main carps and exotic carp like *Hypophthalmichthys molitrix*—have seen notable expansion in India (Jawdhari et al., 2022). Still, disease outbreaks continue to be a constant hazard to aquaculture output, frequently causing significant financial losses and ecological disturbances (Behringer et al., 2020).

Among the etiological agents influencing fish health, bacterial infections are a main source of illness and mortality, especially in populations under environmental stress or intensively grown. Freshwater systems have extensively recorded infections caused by *Aeromonas*, *Pseudomonas*, *Edwardsiella*, and *Flavobacterium* species using a range of virulence techniques including toxin synthesis, tissue-degrading enzymes, and immune evasion mechanisms (Leung et al., 2019, Wise et al., 2021, Quin et al., 2022, Semwal et al., 2023 and Nokhwal et al., 2025). When fish are immunocompromised from environmental or physiological stress, these bacteria could either be opportunistic invaders or main pathogens.

Fish are constantly exposed to diverse microbial communities through their skin, gills, and gastrointestinal system, so their microbial profile quite reflects the nearby aquatic environment (Sehnal et al., 2021). Pathogenic bacteria emit extracellular enzymes including proteases, lipases, lecithinases, and chitinases which help them to destroy host tissues, upset physiological barriers, and colonize interior organs. These enzymatic activities are now recognized as essential virulence determinants since they directly help to start and spread an infection (Sharma et al., 2023 and Lazar et al., 2023).

Often indicating anemia and immunological stress, physiological reactions to bacterial infections show as changes in hematological markers including reduced red blood cell count, hemoglobin concentration, and packed cell volume. Important organs like the liver, kidney, and gills show histopathological alterations suggestive of a systemic infection and tissue destruction (Hinton, 2021). Concurrently with these metabolic changes brought on by infection-induced catabolism of muscle proteins, loss of energy stores, and oxidative stress influencing lipid and vitamin levels, are dietary abnormalities (Wensveen et al., 2024).

An important inland water body sustaining catch fisheries and local livelihoods, Limboti Dam is situated in Ahmedpur taluka of Latur district in Maharashtra. Though important, thorough studies assessing the bacterial illness burden and its effects on fish health in this reservoir are few. *Aeromonas hydrophila* and *Pseudomonas fluorescens* have been found as main fish infections in Marathwada according past regional studies; their exact effects on the physiological and nutritional health of *Hy*pophthalmichthys molitrix have not been methodically studied (Verma, 2023). With this backdrop, the present work aims to close this gap by separating and identifying bacterial pathogens from infected *Hypoph*thalmichthys molitrix in Limboti Dam, assessing their enzymatic pathogenicity, and quantifying the consequent physiological and

nutritional disturbances. The project is to contribute to evidence-based disease surveillance and aquaculture management in inland waters of Maharashtra by creating pathogen profiles and corresponding them with fish health indices.

2. Review of Literature

2.1. Importance of Hypophthalmichthys molitrix in Inland Aquaculture

Rapid growth rate, plankton-feeding behavior, and compatibility with polyculture systems make *Hypophthalmichthys molitrix* among the most important exotic fish species brought to Indian inland aquaculture (Jui et al., 2018). Using phytoplankton makes it environmentally helpful in reducing algal blooms and increasing reservoir biological production. Prakash et al., 2018 reported that because of its non-competitive feeding behavior, which makes it excellent for resource optimization in multi-species systems, *Hypoph*thalmichthys molitrix is rather important in composite fish farming.

2.2. Nutritional Importance of *Hypophthalmichthys molitrix*

Hypophthalmichthys molitrix is a widely cultivated freshwater species known not only for its rapid growth and ecological compatibility but also for its high nutritional value, making it a valuable food fish in many Asian and Eastern European countries (Jawdhari et al., 2022). As a filter-feeding fish, *Hypoph*thalmichthys molitrix efficiently converts phytoplankton and suspended detritus into quality protein, contributing to sustainable aquaculture and public nutrition.

1. High-Quality Protein Source

*Hypo*phthalmichthys molitrix is rich in high-quality, easily digestible protein, making it an important dietary component in protein-deficient regions. On average, the protein content in the edible portion of *Hypo*phthalmichthys molitrix ranges from 17% to 20%, depending on age, diet, and habitat (Fang et la. 2022). These proteins contain all essential amino acids required for human health, including lysine, leucine, methionine, and valine, which are critical for muscle repair, immune function, and enzymatic activity.

2. Balanced Lipid Profile

While relatively low in total fat (approximately 4–8% in the fillet), *Hypo*phthalmichthys molitrix contains a significant proportion of polyunsaturated fatty acids (PUFAs), including omega-3 fatty acids such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) (Djordjevic et al., 2019). These lipids are essential for cardiovascular health, neurological development, and anti-inflammatory responses. Studies have shown that regular consumption of

Hypophthalmichthys molitrix can contribute to reducing the risk of heart disease and improve lipid metabolism (Wang et al., 2023). As *Hypophthalmichthys molitrix* having plenty of health benefits, it can also be beneficiary on post covid complications in human beings.

3. Rich in Vitamins and Minerals

According to Wang et al., 2019, *Hypo*phthalmichthys molitrix is a good source of fat-soluble vitamins such as vitamin A and D, as well as water-soluble B-complex vitamins like B12, niacin, and riboflavin. These vitamins play key roles in vision, skin health, red blood cell formation, and nervous system function. It also provides important **micronutrients** including:

- **Calcium and Phosphorus** – important for bone health
- **Iron** – essential for oxygen transport and hemoglobin synthesis
- **Zinc and Selenium** – antioxidants that support immune and cellular functions

4. Low Caloric Value

Due to its low fat and high water content, *Hypophthalmichthys molitrix* is considered a low-calorie fish, making it suitable for weight management and cardiac patients (Warren, 2020). It is often recommended as a healthy alternative to red meat and high-fat seafood.

5. Affordable Nutritional Security

As a fast-growing species that can be raised in ponds, cages, and reservoirs with minimal feed input, *Hypo*phthalmichthys molitrix offers a cost-effective source of animal protein for rural and urban populations alike (Mohammed et al., 2024). Its production contributes to food security, especially in low-income households and regions with limited access to marine fish.

2.3. Susceptibility of Hypophthalmichthys molitrix to Bacterial Infections

Though resistant to many bacterial diseases, Hypophthalmichthys molitrix is vulnerable in open water systems where environmental factors vary greatly. Particularly in unmonitored reservoirs, bacterial illnesses such motile *Aeromonas* septicemia (MAS), *Pseudomonas*, and columnaris disease are well documented to cause great morbidity and death in *Hypophthalmichthys* molitrix populations (Verma, 2023). Particularly in situations with low water quality and strong organic load, opportunistic bacteria such *Aeromonas hydrophila*, *Pseudomonas fluorescens*, and *Flavobacterium* columnare are routinely recovered from sick fish (Zaheen et al., 2022).

2.4. Pathogenic Mechanisms of Bacterial Isolates

According to Suleiman, 2024, pathogenic bacteria mostly affect tissue invasion and immune suppression by means of virulence factors, therefore enabling their effects:

- **Protease:** Tissue necrosis and weaker epithelial barriers follow from the breakdown of structural proteins by proteases.
- **Lipases:** Lipases cause disturbance of lipid membranes, hence promoting inflammation and cellular lysis.
- **Lecithinases**: Lecithinases also known as phospholipases— hydrolyze phosphatidylcholine in membranes, therefore facilitating bacterial spread.

Though little investigated in fish diseases, chitinases can influence fish gut flora and hence affect digestion indirectly. By neutralizing host defensive systems, these enzymes also help bacterial survival. Under stress in the aquatic environment, studies have found that virulence gene expression—including aerA (aerolysin), hlyA (hemolysin), and lipA (lipase)—is elevated (Sun et al., 2025).

2.5. Environmental Drivers of Pathogen Outbreaks

Conditions favorable for bacterial growth arise from changing water quality characteristics including high ammonia levels, low dissolved oxygen, and varying pH. Like many Indian lakes, Limboti Dam sees seasonal fluctuations and agricultural runoff that could cause eutrophication and immunosuppression in fish. According to Ibáñez et al. (2023), environmental stresses not only raise pathogen virulence but also decrease fish immunity, hence fueling major disease epidemics.

2.6. Haematological Changes in Infected Fish

Perfect bioindicators of pathogen exposure and water quality are fish. Common infections produce:

- ***Anaemia:*** drop in RBC and Hb
- ***Leucocytosis:*** Leukocytosis or leukopenia may occur depending on the stage and severity of infection (Grant, 2015).
- ***Altered PCV, MCHC, and MCV levels***: These changes reflect systemic reactions to infection, including decreased erythropoiesis and poor oxygen transport (McKenzie & Laudicina, 1998 and Mekenye, 2024). Reduced hemoglobin and packed cell volume as observed in *Hy*pophthalmichthys molitrix infected with *Aeromonas* point to chronic illness development (Suhail & Mustafa, 2021).

2.7. Histopathological Impacts of Bacterial Pathogens

Direct visual proof of disease is given by histopathological changes. Important alterations in infected *Hypo*phthalmichthys molitrix consist in:

- Gills: Lamellar fusion, necrosis, and epithelial hyperplasia all of which impede osmoregulation and breathing.
- Liver: Cytoplasmic vacuolization influencing detoxifying; hepatic necrosis; fatty degeneration
- Kidney: Tubular necrosis and glomerular degeneration compromise excretion.

Particularly under environmental stress, these alterations lower physiological resilience and raise mortality risk (Alsudani et al., 2024).

2.8. Nutritional Depletion in Infected Fish

Stress caused by pathogens changes metabolic priority from growth to immunological defense, therefore resulting in:

- ***Protein Catabolism:*** Extended infections break down muscle proteins for use as energy.
- ***Lipid Peroxidation:*** Oxidative stress lowers fat stores, therefore influencing energy and reproductive health.
- ***Micronutrient Loss:*** Under inflammatory conditions, vitamins A, C, E, together with B-complex vitamins, are lost. Reduced market value and customer acceptability follow from this, which emphasizes the need of disease-preventive policies.

2.9. Regional Studies in Maharashtra

There are little methodical research on fish bacterial infections in open reservoirs in Maharashtra. Although Barde, (2023) and Deshmukh et al., (2024) lacked physiological or nutritional data, they reported isolation of *Micrococcus species* and *Pseudomonas fluorescens* from sick *Catla catla* in neighboring reservoirs. Though anecdotal evidence of seasonal fish deaths points to suggesting that silver carp populations in Limboti Dam may require urgent disease monitoring and intervention, none has yet been done. This emphasizes how urgently the present work is needed for data-driven fish health monitoring.

2.10. Global and National Surveillance Gaps

While international agencies like OIE and FAO advocate integrated aquatic health frameworks, India's inland reservoir systems sometimes lack institutional surveillance of fish diseases. Although the Central Institute of Freshwater Aquaculture (CIFA) has underlined the need of

region-specific pathogen profiling to control developing threats, application in semi-natural water bodies like Limboti is still rare.

3. Research Objectives

The present work aims to investigate the contribution of bacterial infections to compromising the physiological and nutritional integrity of *Hypophthalmichthys molitrix* populations in Limboti Dam, Ahmedpur. Investigating the microbial etiology, enzymatic virulence factors, and related host responses will help to produce region-specific data applicable to fish health monitoring and sustainable aquaculture management.

The specific objectives of the study are as follows:

1. **To isolate and identify bacterial pathogens** associated with diseased *Hypo*phthalmichthys molitrix from Limboti Dam using morphological, biochemical, and molecular techniques.
2. **To evaluate the production of extracellular virulence enzymes** (proteases, lipases, lecithinases) by the isolated pathogens and assess their pathogenic potential.
3. **To assess the physiological impact of bacterial infection** through haematological (Hb, RBC, WBC, PCV) and histopathological analysis of critical organs (gills, liver, kidney).
4. **To determine the nutritional alterations in infected fish**, including changes in total protein, lipid, carbohydrate, and vitamin content compared to uninfected individuals.
5. **To study the influence of environmental factors** such as pH, temperature, and salinity on bacterial growth dynamics and virulence expression.
6. **To correlate bacterial load and enzymatic activity** with the extent of physiological and nutritional deterioration in infected fish.

4. Proposed Hypotheses

The study is set around the following assumptions to statistically support the links between host health factors and bacterial infection: Pathogen load and virulence factor expression are favorably linked to statistically significant physiological and nutritional alterations in *Hypo*phthalmichthys molitrix resulting from bacterial infection.

5. Materials and Methods

The study will be carried out applying a multidisciplinary approach including field sampling, microbiological identification, enzymatic profiling, physiological and nutritional analysis, and

evaluation of environmental factors. Selected techniques have been chosen depending on accepted standards in aquatic microbiology and fish pathology.

5.1 Study Area and Sampling Plan

Limboti Dam in Ahmedpur taluka of Latur district, Maharashtra, will house *Hypo*phthalmichthys molitrix specimens. Supporting small-scale and subsistence fishing as well as an annual inland water body, the dam is To accommodate temporal fluctuations in microbial load and environmental conditions, sampling will be conducted seasonally—pre-monsoon, monsoon, post-monsoon.

Fish specimens showing obvious symptoms of bacterial infection—ulcerations, hemorrhages, fin rot, or unusual behavior—will be gathered with gill or cast nets. Healthy-looking specimens will also be collected for comparative analysis as controls.

5.2 Isolation and Enumeration of Bacterial Pathogens

Aseptically swabs of infected external surfaces (skin lesions, fins) and internal tissues (liver, kidney) will be processed for microbiological study.

- Viable Count method: Swabs will be suspended in sterile saline (0.85% NaCl), serially diluted, plated on general (Nutrient Agar) and selective medium (MacConkey Agar), using the spread plate technique. Plates will be incubated for 24 to 48 hours at 28–30°C to ascertain total colony-forming units (CFU/g tissue).
- Pure Culture Isolation: Dominant colonies will be sub-cultured to to obtain axenic cultures for further identification.

5.3 Identification of Bacterial Isolates

Standard microbiological techniques will allow one to identify bacterial isolates:

- Morphological Analysis: Gram staining properties and morphological analysis (shape, size, pigment, edge, elevation) of a colony.
- Biochemical Characterization: Following techniques from Bergey's Manual of Systematic Bacteriology (Krieg and Holt, 1984), tests will include oxidase, catalase, triple sugar iron (TSI) agar, citrate use, urease, indole production, methyl red–Voges-Proskauer (MR–VP), nitrate reduction, and motility assays.

- Molecular Identification: Selected isolates—especially *Aeromonas* and *Pseudomonas* spp.—will be validated by PCR amplification of 16S rRNA genes with genus-specific primers.

5.4 Extracellular Enzyme Assays

Screening for enzyme production will help to evaluate virulence potential:

- Protease Activity: Assessed using skim milk agar; zone of casein hydroly suggests activity.
- Lipase Activity: Clear halo on tributyrin agar shows lipid breakdown.
- Lecithinase Activity: Evaluated on egg yolk agar; opaque precipitation zone around colonies shows lecithinase generation.

If sufficient enzyme production, quantitative experiments can be carried out utilizing spectrophotometric techniques.

5.5 Environmental Parameter Assessment

Sampling site physico-chemical water quality parameters will be tested in situ with portable tools:

- **Temperature and pH:** Digital thermometer and pH meter.
- **Dissolved Oxygen and Ammonia:** Winkler's titration and Nessler's reagent method.
- **Salinity and Conductivity:** Handheld refractometer and conductivity meter

Growing isolates in controlled lab circumstances simulating varying pH (6–9), temperature (20–35°C), and salinity (0.1–1.0%) will allow one to examine effects of these elements on bacterial growth and enzyme production.

5.6 Haematological and Histopathological Studies

- **Blood Collection:** Blood collecting with EDTA-coated syringes from the caudal vein.
- **Parameter Analyzed:** Hemoglobin (cyanmethemoglobin method), RBC and WBC count (haemocytometer), PCV (microhematocrit method), MCH, MCV, and MCHC computed correspondingly.
- **Histopathology:** Tissues (gills, liver, kidney) will be fixed in 10% buffered formalin, processed, embedded in paraffin, sectioned (5 µm), stained with hematoxylin and eosin (H&E), then seen under light microscopy for pathogenic lesions.

5.7 Nutritional Analysis of Fish Tissue

- **Moisture and Ash:** muffle furnace and drying oven gravimetric analysis.
- **Protein Content:** Kjeldahl nitrogen analysis or Lowry technique estimates
- **Lipid Content:** Found by Soxhlet extraction using petroleum ether.

- **Carbohydrates:** Carbohydrates estimated using the Anthrone method..
- **Vitamin Estimation:** If facilities allow, high-performance liquid chromatography (HPLC) will be used for several vitamins (e.g., A, C, E).

Comparatively, samples from sick and healthy fish will be sought statistically significant variations.

5.8 Statistical Analysis

Using suitable statistical program (e.g., SPSS, R, or MS Excel), data will be examined:

- **Descriptive Statistics:** Range, standard deviation, and mean in descriptive statistics.
- **Comparative Analysis:** ANOVA for nutritional and physiological factors independent t-tests
- **Correlation and Regression:** Establish links among fish health measures, enzymatic activity, and bacterial burden.

A significance level of $p < 0.05$ will be considered statistically significant.

6. Current Status

6.1 International Status

Because of its fast development and effective filter-feeding capacity, *Hypo*phthalmichthys molitrix is extensively grown all throughout Asia and Eastern Europe. But bacterial infections have become a major threat, especially in ecologically challenged or intensively farmed settings.

The biggest *Hypo*phthalmichthys molitrix producer, China, has studied bacterial illnesses afflicting this species in great detail (Li et la., 2021). *Aeromonas hydrophila* is a main pathogen generating Motile *Aeromonas* Septicemia (MAS) in Hypophthalmichthys molitrix (Rashid et al., 2013 and Korni et al., 2020). Expressing virulence genes including aerA, hlyA, and act, pathogenic strains has been reported to cause hemorrhages, organ necrosis, and lower nutritional quality of fish muscle.

From sick *Hypo*phthalmichthys molitrix showing liver damage and aberrant haematological profiles, *Aeromonas* veronii and *Pseudomonas* aeruginosa were isolated in Vietnam (Hoai et al. 2019). These infections seriously reduced protein content and changed hemoglobin and RBC levels.

Hwihy et al., 2024 have connected water stress and poor biosecurity to outbreaks of *Edwardsiella* tarda and *Aeromonas* sobria in *Hypophthalmichthys molitrix* , so affecting tissue composition and immunological response.

Rising temperatures, eutrophication, and organic pollution have been underlined by global health agencies such as the OIE and FAO as increasing bacterial virulence specifically affecting fish

metabolism, immunity, and nutrition retention. This has spurred investigation on stress-resistant breeds and probiotics.

Recent proteomics-based studies in China and Hungary have shown that chronic bacterial infections cause catabolism of oxidative lipid damage and depletion of essential vitamins in *Hypophthalmichthys molitrix*, so greatly reducing their market value and nutritional status (Ngan et al., 2024).

6.2 National Status

Although *Hypo*phthalmichthys molitrix is not the principal aquaculture species in India (compared to Indian main carps), *Hypo*phthalmichthys molitrix is extensively included into polyculture systems for their ecological advantages. However, disease surveillance in *Hypophthalmichthys molitrix* remains limited.

Bhubaneswar's ICAR-CIFA has noted how sensitive *Hypo*phthalmichthys molitrix are to *Aeromonas* spp., *Pseudomonas* spp., and *Edwardsiella* spp. under reservoir terms and polyculture (Pathak et al., 2022). These infections have been found by biochemical investigations to lower muscle protein, lipid content, and general flesh quality in infected fish (CIFA Annual Report, 2020–21).

Palaniyappan et al. (2023) reported *A. hydrophila* infection in *Hypophthalmichthys molitrix* from Tamil Nadu fish farms, with haematological alterations suggestive of systemic infection: lowered Hb and RBC levels and raised WBCs. Nutritional study found that muscle tissues' crude protein and vitamin A content had dropped significantly.

Laltlanmawia et al., 2023 found regular isolation of bacterial pathogen from Catla and *Hypo*phthalmichthys molitrix in surrounding reservoirs of Tripuri. But their study concentrated more on identification than on physiological or nutritional influence, therefore highlighting a critical research gap addressed by this study.

Multi-drug resistant *Aeromonas* strains discovered in fish farms in Awankot, Rupnagar, Punjab, India Bengal by Kaur et al. (2024) raise questions regarding not just fish mortality but also the loss of meat quality due to chronic infections that modify nutritional reserves.

Notwithstanding these results, there are lacking systematic national-level investigations linking bacterial infection with nutritional degradation and physiological markers (e.g., histomorphology, blood biochemistry) in *Hypophthalmichthys molitrix*. This emphasizes the originality and need of the suggested study.

7. Expected Outcomes

The present work is expected to produce the following results, therefore supporting field-level application in aquaculture as well as academic knowledge:

1. The research will lead to the isolation and biochemical/molecular identification of major pathogenic bacterial species influencing *Hypophthalmichthys molitrix* in Limboti Dam, so offering fresh understanding of region-specific fish infections.
2. Evaluation of extracellular enzyme production—that of proteases, lipases, and lecithinases— will establish the virulence potential of isolated isolates and hence contribute to our knowledge of their pathophysiology.
3. Haematological and histological impact: The study will measure changes in haematological indices and histological structure in infected fish, therefore producing physiological biomarkers of bacterial disease.
4. A thorough analysis of protein, lipid, carbohydrate, and vitamin levels between infected and uninfected specimens will expose the nutritional toll of chronic bacterial infections.
5. Environmental Influence on Pathogen Behavior: The project will show how abiotic factors control bacterial growth and virulence by modeling environmental variances, thereby enhancing disease risk prediction models.
6. Statistical connections between bacterial load, enzymatic activity, and host health indices will give a quantitative basis for connecting infection degree to physiological and nutritional stress.
7. Future epidemiological research, risk-reducing initiatives, and health management measures in open reservoir fisheries systems will find reference in the data produced here.

8. Significance of the Study

This study has great scientific, financial, and environmental value as well:

- **Scientific Advancement:** It will close a significant knowledge gap in fish microbiology and aquatic pathology particular to Marathwada's inland reservoirs. Unlike regulated aquaculture systems, wild reservoirs show intricate ecological interactions that, in the framework of pathogen dynamics, remain under investigated.
- **Aquaculture Health Management:** Identification of important infections and their health consequences will help to design early disease detection systems and maybe advise preventative or therapeutic approaches catered for the area.
- **Food Safety and Nutritional Security:** The study tackles consumer-level issues by means of an analysis of how bacterial illnesses affect the nutritional value of fish, therefore tying environmental microbial dangers to food quality and human nutrition.
- **Policy and Resource Planning:** The results can help local fisheries departments and reservoir management authorities create surveillance and intervention plans meant to reduce disease outbreaks and maintain fish output.

- **Regional Research Value:** To date, very few systematic investigations have been carried out in Limboti Dam or nearby districts, hence this work is a pioneering effort that might inspire more research projects and monitoring activities all across Maharashtra.

8. Work Plan and Timeline

Year	Key Activities
Year 1 2025-26	- Completion of PhD coursework - Extensive literature review and refinement of research objectives
	- Investigation and selection of sampling sites in Limboti Dam - Baseline environmental monitoring (pH, temperature, salinity, DO, ammonia) - Initial fish sampling and isolation of bacterial pathogens
Year 2 2026-27	- Morphological and biochemical identification of isolates - Molecular confirmation (16S rRNA PCR)
	- Screening of virulent enzymes (protease, lipase, lecithinase) - Growth studies under varying environmental conditions (pH, salinity, temperature)
Year 3 2027-28	- Haematological analysis (Hb, RBC, WBC, PCV) - Histopathology of gills, liver, kidney
	- Nutritional analysis (protein, lipids, carbohydrates, vitamins) - Statistical comparison: healthy vs. infected fish
Year 4 2028-29	- Integration and interpretation of results - Drafting of thesis chapters
	- Compilation of graphs, histological images, and statistical outputs - Final thesis submission and Viva-voce.

PhD Work Tentative Plan - Gantt Chart (2025–2029)



References

Alsudani, H. M., Al-Shammari, N. A., & Al-Niaeem, K. S. (2024). Histopathological alterations of intestines, liver, kidneys, and spleen in Common carp, Cyprinus carpio L. infected by Staphylococcus lentus. *International Journal of Aquatic Biology*, *12*(6), 563-568.

Barde, R. D. (2023). Exploring the Disease-Causing Potential of Aeromonas Hydrophila Obtained from Freshwater Carp-Cirrhinus mrigala. *J. Surv. Fish. Sci*, *10*, 1725-1732.

Behringer, D. C., Wood, C. L., Krkošek, M., & Bushek, D. (2020). Disease in fisheries and aquaculture. *Marine disease ecology*, *183*.

Deshmukh, A. D., Jadhav, M. P., Achegawe, R. M., & Puri, D. G. (2024). Drug-resistant Bacterial Pathogens: Isolation and characterization in freshwater fish *Catla catla*. *International Journal of Fisheries and Aquatic Research:* 9, 1,16-21

Djordjevic, V., Trbovic, D., Nastasijevic, I., Ljubojevic, D., Lakicevic, B., & Nikolic, A. (2019). Effect of frying on the fatty acid composition of silver carp and common carp. *Scientific journal" Meat Technology"*, *60*(1), 30-36.

Fang, D. A., Sun, H., Peng, Y., Kuang, Z., Zhou, Y., & Xu, D. (2022). Living status and perspective of the Silver Carp (Hypophthalmichthys molitrix) in the lower reach of the Yangtze River: Insights from population distribution, age structure, and habitat preference analyses. *Fishes*, *7*(5), 254.

Grant, K. R. (2015). Fish hematology and associated disorders. *Veterinary Clinics: Exotic Animal Practice*, *18*(1), 83-103.

Hinton, D. E. (2021). Toxicologic histopathology of fishes: a systemic approach and overview. In *Pathobiology of marine and estuarine organisms* (pp. 177-215). CRC Press.

Hoai, T. D., Trang, T. T., Van Tuyen, N., Giang, N. T. H., & Van Van, K. (2019). Aeromonas veronii caused disease and mortality in channel catfish in Vietnam. *Aquaculture*, *513*, 734425.

Hwihy, H. M., Zeina, A. F., Husein, M. S. A., & El-Tabakh, M. A. (2024). Biofloc biosecurity: A revolutionary paradigm in augmenting aquaculture health and resilience against Aeromonas hydrophila. *Aquaculture and Fisheries*.

Ibáñez, A., Garrido-Chamorro, S., & Barreiro, C. (2023). Microorganisms and climate change: a not so invisible effect. *Microbiology Research*, *14*(3), 918-947.

Jawdhari A, Mihăilescu DF, Fendrihan S, Jujea V, Stoilov-Linu V, Negrea BM. Silver Carp (*Hypophthalmichthys molitrix*) (Asian Silver Carp) Presence in Danube Delta and Romania-A Review with Data on Natural Reproduction. Life (Basel). 2022 Oct 12;12(10):1582. doi: 10.3390/life12101582. PMID: 36295017; PMCID: PMC9605455.

Jui, R., Haque, M. & Rahmatullah, S. M. (2018). Growth performance of silver carp (Hypophthalmichthys molitrix) in cage stocked at different densities. Journal of the Bangladesh Agricultural University. 16. 322-327. 10.3329/jbau.v16i2.37989.

Kaur, S., Kaur, H., Kaur, B., Kumar, B. N., Tyagi, A., Singh, P., ... & Munang'andu, H. M. (2024). Isolating pathogenic multidrug-resistant Aeromonas hydrophila from diseased fish and assessing the effectiveness of a novel lytic Aeromonas veronii bacteriophage (AVP3) for biocontrol. *Microbial Pathogenesis*, *196*, 106914.

Korni, F. M., Abo El-Ela, F. I., & Moawad, U. K. (2020). Role of Moringa oleifera leaves and aqueous extract in prevention of Motile Aeromonas Septicemia in common carp, Cyprinus carpio fingerlings with a reference to histopathological alterations. *Aquaculture International*, *28*, 153-168.

Laltlanmawia, C., Saha, H., Ghosh, L., Saha, R. K., & Malla, S. (2023). Identification and analysis of pathogenic bacteria causing outbreaks in Indian major carp aquaculture of Tripura. *Journal of Aquatic Animal Health*, *35*(4), 263-279.

Lazar V, Oprea E, Ditu LM. Resistance, Tolerance, Virulence and Bacterial Pathogen Fitness-Current State and Envisioned Solutions for the Near Future. Pathogens. 2023 May 22;12(5):746. doi: 10.3390/pathogens12050746. PMID: 37242416; PMCID: PMC10222143.

Leung KY, Wang Q, Yang Z, Siame BA. *Edwardsiella piscicida*: A versatile emerging pathogen of fish. Virulence. 2019 Dec;10(1):555-567. doi: 10.1080/21505594.2019.1621648. PMID: 31122125; PMCID: PMC6592360.

Li, D., Prinyawiwatkul, W., Tan, Y., Luo, Y., & Hong, H. (2021). Asian carp: A threat to American lakes, a feast on Chinese tables. *Comprehensive Reviews in Food Science and Food Safety*, *20*(3), 2968-2990.

Little, D. C., Newton, R. W., & Beveridge, M. C. M. (2016). Aquaculture: a rapidly growing and significant source of sustainable food? Status, transitions and potential. *Proceedings of the Nutrition Society*, *75*(3), 274-286.

McKenzie, S. B., & Laudicina, R. J. (1998). Hematologic changes associated with infection. *Clinical Laboratory Science*, *11*(4), 239.

Mekenye, O. J. (2024). *Profile Of Haematological Indices Among Pulmonary Tuberculosis Patients Attending Kisii Teaching And Referral Hospital, Kisii Kenya* (Doctoral dissertation, Kisii University).

Mohammed HH, Ebrahim M, Youssef MI, Saleem AY, Abdelkhalek A. Behavior and management of carp fish: A review. Open Vet J. 2024 Jan;14(1):1-11. doi: 10.5455/OVJ.2024.v14.i1.1. Epub 2024 Jan 31. PMID: 38633176; PMCID: PMC11018436.

Ngan BNM, Heyse J, Delamare-Deboutteville J, Props R and Shelley CC. (2023). A review of microbiomes in carp polyculture systems. Penang, Malaysia: WorldFish. Technical Report: 2023-06.

Nokhwal, A., Vaid, R. K., Anand, T., Verma, R., & Gulati, R. (2025). *Aeromonas* Species Diversity, Virulence Characteristics, and Antimicrobial Susceptibility Patterns in Village Freshwater Aquaculture Ponds in North India. *Antibiotics*, *14*(3), 294. https://doi.org/10.3390/antibiotics14030294

Palaniyappan, S., Sridhar, A., Kari, Z. A., Téllez-Isaías, G., & Ramasamy, T. (2023). Potentials of Aloe barbadensis inclusion in fish feeds on resilience to Aeromonas hydrophila infection in freshwater fish Labeo rohita. *Fish Physiology and Biochemistry*, *49*(6), 1435-1459.

Pathak, H., Mishra, J. P., & Mohapatra, T. (2022). Indian agriculture after independence. *Indian Council of Agricultural Research, New Delhi*, *110*(001), 426.

Prakash, B., Khairnar, S., Mandal, A. Kumar, A. & Kumar, B. (2018). Composite fish farming: A review on economic enterprise for rural empowerment and livelihood generation.

Qin S, Xiao W, Zhou C, Pu Q, Deng X, Lan L, Liang H, Song X, Wu M. Pseudomonas aeruginosa: pathogenesis, virulence factors, antibiotic resistance, interaction with host, technology advances and emerging therapeutics. Signal Transduct Target Ther. 2022 Jun 25;7(1):199. doi: 10.1038/s41392-022-01056-1. PMID: 35752612; PMCID: PMC9233671.

Rashid, M., Hossain, M. S., & Ali, M. F. (2013). Isolation and identification of Aeromonas hydrophila from silver carp and its culture environment from Mymensingh region. *Journal of the Bangladesh Agricultural University*, *11*(2), 373-376.

Sehnal, L., Brammer-Robbins, E., Wormington, A. M., Blaha, L., Bisesi, J., Larkin, I., ... & Adamovsky, O. (2021). Microbiome composition and function in aquatic vertebrates: small organisms making big impacts on aquatic animal health. *Frontiers in microbiology*, *12*, 567408.

Semwal A, Kumar A, Kumar N. A review on pathogenicity of *Aeromonas hydrophila* and their mitigation through medicinal herbs in aquaculture. Heliyon. 2023 Feb 28;9(3):e14088. doi: 10.1016/j.heliyon.2023.e14088. PMID: 36938468; PMCID: PMC10018484.

Sharma, I., & Yumnam, H. (2023). Introduction to Microorganisms and Pathogenesis. In *Antimicrobial Photodynamic Therapy* (pp. 1-17). CRC Press.

Suhail, R. N., & Mustafa, S. A. (2021). Antibacterial Effect of Silver Nanoparticles on Hematological and Biochemical Parameters in Common Carp Infected with Aeromonas hydrophila. *Annals of the Romanian Society for Cell Biology*, *25*(6), 5074-5086.

Suleiman, R. Z. (2024). A comparative approach to understanding bacterial pathogenicity: The role of enzymatic and hemolytic virulence factors. *Journal of Medical Genetics and Clinical Biology*, *1*(6), 150-161.

Sun, Y., Hao, Y., Wang, Z., Wang, H., & Gao, Y. (2025). Isolation, characterization, and whole genome sequencing analysis of Aeromonas veronii from Channa argus in China. *Veterinary Research Communications*, *49*(1), 1-18.

Verma, A. K. (2023). *Studies on bacterial diseases of cultured freshwater fishes* (Doctoral dissertation, Chaudhary Charan Singh Haryana Agricultural University).

Wang AnLi, W. A., Zu JinFeng, Z. J., Shi WenQiang, S. W., Tan ChengFeng, T. C., Li XiangLan, L. X., Lu JianFeng, L. J., & Lin Lin, L. L. (2019). Analysis and evaluation of the nutritional components in swim bladder of silver carp.

Wang, Q., Wang, R., Zhao, X., Lu, H., Zhang, P., Dong, X., & Wang, Y. (2023). Comparison of the effect of phospholipid extracts from salmon and silver carp heads on high-fat-diet-induced metabolic syndrome in C57BL/6J Mice. *Marine Drugs*, *21*(7), 409.

Warren, D. C. (2020). *Assessment of Protein Quality, Lipid Metabolism, Hepatic Transcriptome and Safety of a Silver Carp (Hypophthalmichthys molitrix) Derived Sarcoplasmic Protein Powder*. West Virginia University.

Wensveen, F. M., Šestan, M., & Polić, B. (2024). The immunology of sickness metabolism. *Cellular & Molecular Immunology*, *21*(9), 1051-1065.

Wise, A. L., LaFrentz, B. R., Kelly, A. M., Khoo, L. H., Xu, T., Liles, M. R., & Bruce, T. J. (2021). A Review of Bacterial Co-Infections in Farmed Catfish: Components, Diagnostics, and Treatment Directions. *Animals*, *11*(11), 3240. https://doi.org/10.3390/ani11113240

Zaheen, Z., War, A. F., Ali, S., Yatoo, A. M., Ali, M. N., Ahmad, S. B., ... & Paray, B. A. (2022). Common bacterial infections affecting freshwater fish fauna and impact of pollution and water quality characteristics on bacterial pathogenicity. In *Bacterial fish diseases* (pp. 133-154). Academic Press.

Research Scholar	**Research Guide**
Baldawa Mohini Rajkumar	**Dr. R. D. Barde**